
DIE EILE_COPY AD



ADJUSTMENT AND VALIDATION OF THE MATHEMATICAL PREDICTION MODEL FOR SWEAT RATE, HEART RATE AND BODY TEMPERATURE
UNDER OUTDOOR CONDITIONS

ANNUAL REPORT

- Y. SHAPIRO
- R. BURSTEIN
- Y. EPSTEIN



NOVEMBER 1, 1986

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND Fort Detrick, Frederick, Maryland 21701-5012

Grant No. DAMD17-85-G-5044

Ministry of Defense
Directorate of Defense R&D, Israel

Approved for public release; distribution unlimited

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

REPORT DOCUMENTATIO	Form Approved OM8 No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION UNGLASSIFIED	16. RESTRICTIVE MARKINGS	
20. SECURITY CLASSIFICATION AUTHORITY	3. DISTRIBUTION / AVAILABILITY OF REPORT	
26. DECLASSIFICATION / DOWNGRADING SCHEDULE	Approved for public release distribution unlimited	2 ;
4. PERFORMING ORCANIZATION REPORT NUMBER(S)	S. MONITORING ORGANIZATION REPORT NU	MBER(S)
66. NAME OF PERFORMING ORGANIZATION (If applicable) Ministry of Defense	79. NAME OF MONITORING ORGANIZATION	
6c ADDRESS (City, State, and ZIP Code) Directorate of Defense R&D, Israel	7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Medical (if applicable) Research & Development Command SGRU-RMI-S	9. PROCUREMENT INSTRUMENT IDENTIFICATI DAMD1 7-85-G-5044	ON NUMBER
Research & Development Command SGRD-RMI-S 8c. ADDRESS (City, State, and ZIP Code)	10. SOURCE OF FUNDING NUMBERS	
Fort Detrick Frederick, Maryland 21701-5012	PROGRAM PROJECT TASK NO. 3E1- NO. 62777A 62777A879 BJ	WORK UNIT ACCESSION NO. 081
11. TITLE (include Security Classification) (U) Adjustment and N Model for Sweat Rate, Heart Rate and Body Ten 12. PERSONAL AUTHOR(S)	Validation of the Mathematical sperature Under Outdoor Condition	
Shapiro, Y., R. Burstein, Y. Epstein		
13b. TYPE OF REPORT Annual 12b. TIME COVERED FROM 10/7/85 TO 10/6/86	14. DATE OF REPORT (Year, Month, Day) 15. 1986 November 1	PAGE COUNT 16
16. SUPPLEMENTARY NOTATION		
	Continue on reverse if necessary and identify b	•
	ody temperature, heat stress, o del, protective garments, heat	
12 01		
Commence of the state of the st	which were developed in USARIE conmental conditions. Twenty matric conditions, solar loads, sysiological parameters were callion models, which were developmentate the physiological results a factor related to long waves some will be investigated during	M, to predict hale subjects clothing refully ed from ponses under radiations
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED SAME AS RPT. DTIC USERS \$23 NAME OF RESPONSIBLE INDIVIDUAL	21. ABSTRACT SECURITY CLASSIFICATION Unclassified 22b TELEPHONE (Include Area Code) 22c. OFF	ICE SYMBOL
Mary Frances Bostian	301/663-7325 SGR	D-RMI-S

COS LA SECRETARIO DE CONTROL DE COSTO DE CONTROL DE CON

П	T				
ы	11				
_	1.3				

ADJUSTMENT AND VALIDATION OF THE MATHEMATICAL PREDICTION MODEL FOR SWEAT RATE, HEART RATE AND BODY TEMPERATURE UNDER OUTDOOR CONDITIONS

ANNUAL REPORT

Y. Shapiro

R. Burstein

Y. Epstein

November 1, 1986

Supported by

U.S. Army Medical Research And Development Command Fort Detrick, Frederick, Maryland 21701-5012

Grant No. DAMD17-85-G-5044

Ministry of Defense Directorate of Defense R&D, Israel

Approved for public release. Distribution unlimited.

The findings in this report are not to be considered as an offical Israel Defense Forces or Department of the Army position unless so designated by other authorized documents.

SUMMARY

During the last two decades the Military Ergonomics Division the U.S. Army Research Institute of Environmental Medicine of have developed a series of predictive equations for physiological responses of clothed soldiers performing physical work in various environmental extremes. However, these predicting models were developed under controlled laboratory conditions. The present study is undertaken to validate these predicting models under butdoor conditions. Twenty fit subjects were exposed to 12 combinations of climate, clothing and metabolic rates. During the experimental sessions physiological and meteorological parameters were collected. The data obtained for rectal temperature sweat rate was compared with the predicted values. It observed that under the autdoor conditions the indoor predicting models for rectal temperature and sweat loss overestimate outdoor actual values. It has been concluded to continue study in an attempt to investigate the outdoor factor which cause a deviation from the predicted values.

Accesi	on For	1			
NTIS DTIC	CRA&I TAB	b D			
Unannounced [] Justification					
By Distribution /					
Availability Codes					
Dist	Avail and Specie				
Ĥ-1		**************************************			



FOREWORD

The objectives of this grant are: a) to evaluate the physiological responses to physical activities at various metabolic levels, performed under different outdoor conditions when different clothing combinations are applied; b) adjust the mathematical models which were developed in USARIEM to predict rectal temperature and sweat rate production for natural outdoor conditions.

The mathematical models for the prediction of rectal temperature and sweat production were developed and assessed under controlled laboratory conditions. The present study is based on the hypothesis that, they will be valid for outdoor conditions with only minor modifications and adjustments.

For the protection of human subjects the investigators have adhered to policies of applicable Federal Law 45CFR46.

Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

ᡮᡱᡠᢜᡠᢢᢆᡡᡠᡳᡭᡳᢥ᠈ᡬᢌŶᢣᡐċᢐᡶᢐᡶᢐᡟᡠᢨᡠᢐᡠᢨᡠᢣ᠈ᠫᢨᡡᢛᢨᢐᠫᢐᢨᡑᡱᢐᢗᢌᡧᡠᡳᡷᡠᡧ᠘ᢞᡠᡧᡠᡐᢖᢒᡈᡐᡐᢠᢒᢠᢒᡑᢒᡑᠪᡑᠪᡑᠪᡑ*ᡡᢎᠽᢐ*ᡊᡑᢐᡑᢐ

TABLE OF CONTENTS

Su mma ry	1
Foreword	2
Table of Contents	3
Introduction	4-5
Material & Methods	5-7
Results	8-12
References	13
Distribution list	14

INTRODUCTION

Three major physiological factors determine the ability of men to tolerate heat while operating under hot climate: body temperature, cardiovascular adjustments and water balance. During the last two decades mathematical models for predicting body temperature, heart rate and sweat rate response to a given heat load, have been developed (1-6). These models is have been validated for numerous combinations of environmental conditions, physical activity and clothing, all under controlled laboratory conditions.

Thermoregulatory balance of the body is determined by the relation between the evaporative cooling required (E) and the maximal evaporative heat loss to the environment (E). Sweat loss is expressed by the same terms:

-0.455 -2 -1 m = 27.9 E E ; (g m h)

E is constituted of the metabolic heat production, heat req transfer characteristics of clothing, and ambient climate conditions; E is dictated by the vapor transfer properties of clothing and the vapor pressure gradient between the skin and the air. In a similar way T and HR are predicted.

Solar radiation which has an additive effect on the thermuregulatory balance is not taken into consideration in these equations. Thus, these models are effective and accourate for indoor conditions, but not necessarily so for outdoor climate. Recent preliminary experiments, using artificial indoor solar radiation by infrared electrical lamps, suggest that the radiant

heat load H should be added to the physical heat exchanged between the body and the environment (E) when the mathematical formula for sweat rate prediction is applied. According to this modification H = K (MRT) when: K is a coefficient which depends on the ratio direct to diffused radiation and on clothing, and MRT is the additive value to mean radiant temperature of solar radiation. MRT can be expressed also as:

MRT =
$$(1 + 2.2 \text{ V})$$
 (T - T); (C)

Where T is the black globe temperature and T is the ambient g a temperature.

In order to apply the predicting models for outdoor conditions, field research was designed.

MATHERIAL AND METHODS

The first stage of the study was conducted during the summer of 1986, under natural climatic conditions of a desert (hot/dry) and coastal (hot/wet) areas of Israel.

<u>Subjects</u>. Twenty fit male volunteers, 17-19 years old participated in the study after being briefed by a senior investigator and signing an informed consent form. The subjects were inhabitants of either the desert or the coastal region, naturally acclimatized to the local climatic conditions.

<u>Preliminary Screening</u>. Prior to experimental testing each subject underwent medical examination including resting E.C.G.,

urinalysis and SMA-12 screening chemistry. To determine aerobic power, oxygen uptake at maximal exercise was analyzed using a computerized metabolic chart (MMC-Horizon; Sensor Medics). A progressive treadmill running test was performed at a constant speed of 8 mph and a stepwise grade increment of 2% every 2 min. until exhausion. The test was conducted indoors under comfort thermal conditions (20 C; 50% relative humidity).

Experimental Design. Two groups of 10 subjects each, were exposed to either hot/dry (desert) or hot/wet (coast) climate. Each subject in either group participated in 12 experimental sessions, assigned to him et random. All experimental sessions were carried out between 11:30-13:30. The experimental combinations were as follows:

- Solar radiation: Sun (maximal solar radiation)
 Shade (minimal solar radiation)
- 2. Clothing: Standard fatigues

CB protective overgarments (MOPP IV)

3. Metabolic rate: Rest (100 watt)

Moderate work (320 watts)

Heavy work (425 watts)

Exercise consisted of treadmill walking at speeds of 4.8 km h -1 and 7.2 km h to elicit the desired average metabolic rate. Each experimental session lasted 120 min., consisted of two bouts of 10 min. of rest and 50 min. of exercise.

<u>Physiological Status Assessment</u>. Prior to exposure and at 5 min. intervals during the test until its end, heart rate and rectal temperature were monitored. Skin temperature was monitored at 15 min intervals. Rectal temperature was recorded with a

Yellow Spring Instrument (YSI) rectal termistor probe inserted 10 cm beyond the anal sphincter. Skin temperature was monitored with a YSI skin termistor. Skin temperature was measured simultanously at 3 locations (chest, calf and forearm). Weighted skin temperature (T) was calculated according to Burton (7). Oxygen consumption was measured during the rest and exercise periods using an automated metabolic chart (MMC-Horizon, Sensor Medics). E.C.G. electrodes were affixed to the subjects chest (C:1-5 placement) and heart rate was recorded parallel to metabolic measurement. Body weight was measured with electronic scale (accurate to ± 10 g). Ad libitum drinking was Water intake and urine output were precisely encouraged. measured. Sweat rate was determined from weight loss (by measuring the subjects weight pre and post exposure) adjusted for water intake and urine output.

<u>Meteorological Parameters</u>. During the experimental sessions the following environmental variables were continuously monitored:

- 1. dry bulb temperature, with a sling psychrometer.
- 2. wet bulb temperature, with a sling psychrometer.
- 3. wind speed, with a cap anamometer.
- 4. black globe temperature, with a 6" copper globe.
- 5. solar radiation, direct radiation and infrared radiation with a Spectral Pyranometer (model PSP, EPLAB U.S.A.).

<u>Calculations</u>. The actual results were compared to the pedicted values using the mathematical models as summarized by Pandolf et al. (8).

RESULTS

The mean ambient temperature in the desert area was slightly o higher than that of the coastal region (32 C and 30 C, respectively). Solar radiation was also similar in both regions -2 (800 watt m). However, relative humidity was about 15% higher in the coastal area than in the desert area (60% and 45%, respectively).

Tables 1 and 2 present the mean rectal temperature after 60 and 120 min. of exposure to ambient climate at the coastal (Table 1) and desert (Table 2) areas. The data is presented in comparison to the predicted values. Table 3 presents the results concerning sweat loss (presented as g h) under the different combinations both for the desert and coastal areas. Consistently, the predicted values, as calculated by the prediction models, are higher than the observations. It is evident that the prediction models overestimate the actual values under all environmental, clothing and metabolic rate combinations by a constant value.

At present, the data collected are analyzed in an attempt to adjust the prediction models for outdoor conditions.

Table 1 - MEAN RECTAL TEMPERATURE (°C) AFTER 60 AND 120 MIN.

OF EXPOSURE, MEASURED (M) V5. PREDICTED (P) UNDER

COASTAL AREA CONDITIONS (30°C, 60%RH)

A. Rest (110 watt)

60 min.

120 min.

		Fat.	MOPP4	Fat.	MOPP4
Shade	М	36.90	37.00	37.10	37.10
	P	37.57	37.38	37.98	37.64
Sun	М	37.20	37.40	37.20	37.50
	P	37.55	37.63	37.95	38.08

B. Moderate exercise (320 watt)

60 min.

120 min.

	Fat.	MOPP4	Fat.	MOPP4
М	37.60	37.90	37.60	38.00
P	37.95	38.19	38.31	38.75
М	37. <i>7</i> 5	38.30	37.80	38.70
P	38.26	38.50	38.90	39.43
	P M	M 37.60 P 37.95 M 37.75	M 37.60 37.90 P 37.95 38.19 M 37.75 38.30	M 37.60 37.90 37.60 P 37.95 38.19 38.31 M 37.75 38.30 37.80

C. Heavy exercise (425 watt)

60 min.

120 min.

		Fat.	MOPP4	Fat.	MUPP4
Shade	М	37.98	38.67	38.00	38.70
	P	38.32	38.72	38.86	39.70
Sun	M	38.19	38.82	38.15	- *
	P	38.61	39.05	39.47	40.53

^{*} could not be sustained within safty limits.

Table 2 - MEAN RECTAL TEMPERATURE (°C) AFTER 60 AND

120 MIN. OF EXPOSURE, MEASURED (M) VS. PREDICTED

(P) UNDER DESERT CONDITIONS (31°C, 45%RG)

A. Rest (110 watt)

~~	
HI I	
au	mın.

120 min.

		Fat.	MUPP4	Fat.	MOPP4
Shade	М	36.80	36.90	36.80	36.90
	P	37.18	37.32	37.29	37.52
Sun	M	37.10	37.00	37.10	37.10
	P	37.40	37.50	37.67	37.85

B. Moderate exercise (320 watt)

60 min.

120 min.

		Fat.	MOPP4	Fat.	MOPP4
Shade	М	37.30	37.30	37.40	37.50
	P	37.83	38.09	38.10	38.54
Sun	М	37.40	37.70	37.50	38.10
	P	38.07	38.34	38.53	39.05

C. Heavy exercise (425 watt)

60 min.

120 min.

		Fat.	MOPP4	Fat.	MOPP4	
Shade	М	37.50	37.60	37.60	37.90	
	P	38.14	38.46	38.53	39,11	
Sun	M	37.60	38.10	37.7 0	38.60	
	P	38.38	38.71	38.98	39.67	

Table 3 - SWEAT LOSS (g h) DURING THE EXPOSURE

A. Rest (110 watt)

		Desert		Coast			
		Fat.	MOPP4	Fat.	MOPP4		
Chada	М	139	264	200	300		
Shade	P	180	270	180	360		
Sun	M	310	440	460	640		
	P	540	630	760	810		

B. Moderate exercise (320 watt)

		Desert		Coast		
		Fat.	MOPP4	Fat.	MOPP4	
Shade	М	410	560	560	790	
	P	810	1080	900	1170	
Sun	M	590	860	760	990	
	P	1080	1350	1260	1530	

C. Heavy exercise (425 watt)

		Des	Desert		Coast	
		Fat.	MOPP4	Fat.	MOPP4	
Shade	М	490	750	790	390	_
	P	1080	1350	1260	1620	
Sun	М	680	1160	960	1490(?)	
	P	1350	1620	1530	1890	

^(?) see note in table 1C.

Experimental layout for the 2nd phase of the study

The data obtained so far was discussed with the American collaborators from the Military Ergonomics Division at the U.S. Army Research Institute of Environmental Medicine. It was concluded that the study should be continued according to the following guidelines:

- 1. Metabolic rate: heavy exercise (425 watt)
- 2. Clothing: standard fatigues
- 3. Environmental conditions: solar load

shade

The study will be carried out both indoors and outdoors. Environmental conditions for indoors studies will simulate outdoor conditions concerning ambient temperature and relative humidity. In addition the possibility to validate the prediction models under extremely hot and very dry ambient conditions (40 C, 20% rh) is considered.

This will enable us to investigate the possibility that for outdoor condition a constant correcting factor should be added to the mathematical models. It is hypothesized that this factor might be related to the solar load, which could not be simulated under laboratory conditions.

REFERENCES

- 1. Givoni B., Goldman RF: Predicting metabolic energy cost. J. Appl. Physiol. 30: 429-433, 1971.
- Givoni B., Goldman RF: Predicting rectal temperature response to work environment and clothing. J. Appl. Physiol. 32: 812-822, 1972.
- Givoni B., Goldman RF: Predicting heart rate response to work, environment and clothing. J. Appl. Physiol. 34: 201-204, 1973.
- Givoni B., Goldman RF: Predicting effects of heat acclimatization on heart rate and rectal temperature.
 J. Appl. Physiol. 35: 875-879, 1973.
- 5. Pandolf KB., Givoni B., Goldman RF: Predicting energy expenditure with loads while standing or walking very slowly. J. Appl. Physiol. 43: 477-581, 1977.
- 6. Shapiro Y., Pandolf KB., Breckenridge JR., Goldman RF:

 Predicting sweat loss response to exercise, environment
 and clothing. Europ. J. Appl. Physiol. 48: 83-96, 1982.
- 7. Burton AC: Human Calorimetry II. The average temperature of the tissue of the body. J. Nutr. 9: 261-280, 1935.
- 8. Pandolf KB., Stroschein LA., Drolet LL., Gonzalez RR., Sawka
 MN: Prediction modeling of physiological responses and
 human performance in the heat. Comp. Biol. Med. 1987,
 in press.

DISTRIBUTION LIST

1	Commander						
	US Army Medical Research and Development Command						
	ATTN: SGRD-RMI-S						
	Fort Detrick, Frederick, MD 21701-5012						

- Commander
 US Army Medical Research Acquisition Activity
 ATTN: SGRD-RMA-RC
 Fort Detrick, Frederick, MD 21701-5014
- Commander
 US Army Medical Research Institute
 of Environmental Medicine (USARIEM)
 ATTN: Dr. Kent Pandolf
 Bldg. 42
 Natick, Massachusetts 01760-5007
- 12 Administrator
 Defense Technical Information Center
 ATTN: DTIC-DDA
 Cameron Station
 Alexandria, Virginia 22314
- Commandant
 Academy of Health Sciences, US Army
 ATTN: AHS-CDM
 Fort Sam Houston, Texas 78234